

April 1997
 SISSA 59/97/EP
 solv-int/9705001

Hamiltonian structure and coset construction of the supersymmetric extensions of $N = 2$ KdV hierarchy

L. Bonora^{*a} and S. Krivonos^{†b}

^(a) *International School for Advanced Studies (SISSA/ISAS)*

Via Beirut 2, 34014 Trieste, Italy

INFN, Sezione di Trieste

^(b) *Bogoliubov Laboratory of Theoretical Physics, JINR,*

141980 Dubna, Moscow Region, Russia

Abstract

A manifestly $N = 2$ supersymmetric coset formalism is applied to analyse the "fermionic" extensions of $N = 2$ $a = 4$ and $a = -2$ KdV hierarchies. Both these hierarchies can be obtained from a manifest $N = 2$ coset construction. This coset is defined as the quotient of some local but non-linear superalgebra by a $\widehat{U(1)}$ subalgebra. Three superextensions of $N = 2$ KdV hierarchy are proposed, among which one seems to be entirely new.

^{*}E-mail: bonora@frodo.he.sissa.it

[†]E-mail: krivonos@thsun1.jinr.dubna.su

1 Introduction

The search for $N = 2$ and $N = 4$ extensions of known hierarchies of integrable equations, has brought new impulse to the research in the field of integrable systems. Not only are these extensions characterized by beautiful supersymmetric second hamiltonian structures, but, what is more important, they seem to be endowed with an organizing capacity, that allows them to encompass different cases under a unique structure. Nevertheless, in many cases adding some additional fields or even changing the statistics of the fields into the game – which are typical ways of getting new hierarchies from old ones – may preserve integrability but make the second Hamiltonian structure non-local [1, 2]. The purpose of this letter is to demonstrate on the simplest possible examples that the systems with non-local second Hamiltonian structure are subsystems of extended integrable systems with a *local* second Hamiltonian structure and can be obtained from the latter via the coset approach. One can hope that this represents the general situation and that the fundamental integrable hierarchies possess local second Hamiltonian structures.

We start from “fermionic” extensions of $N = 2, a = 4$ [3, 1] and $N = 2, a = -2$ [4] KdV hierarchies which possess the same non-local second hamiltonian structure. It turns out that the best way to unify these hierarchies is via a coset construction which ‘localizes’ the second hamiltonian structure. We explicitly construct the corresponding coset in section 2. In section 3 we show that the Dirac reduction applied to the corresponding local but non-linear algebra reproduces the non-local algebra that constitutes the original second Hamiltonian structure. In section 4, as the first step to analyse these extended systems, we search for integrable systems that have a subalgebra of our non-linear algebra as their second Hamiltonian structure. Two solutions correspond to supersymmetric extensions of already known integrable hierarchies, but we also introduce the first flows and hamiltonians of an $N = 2$ hierarchy, which does not seem to have any correspondence among the cases studied so far.

2 ”Fermionic” extensions of the N=2 KdV in the coset approach

2.1 Preliminaries: N=4 KdV hierarchy

The $N = 4$ supersymmetric KdV equation has been constructed in [5]. In terms of $N = 2$ superfields it can be written as the coupled system of equations for the general bosonic superfield J and for a pair of chiral/anti-chiral bosonic superfields $\Phi/\bar{\Phi}$. Explicitly, the first two non-trivial flows (second and third) have the following form

$$\begin{aligned} \frac{\partial J}{\partial t_2} &= -[\mathcal{D}, \bar{\mathcal{D}}] J' - 4JJ' - (\Phi\bar{\Phi})', \\ \frac{\partial \Phi}{\partial t_2} &= -\Phi'' + 4\mathcal{D}\bar{\mathcal{D}}(J\Phi), \quad \frac{\partial \bar{\Phi}}{\partial t_2} = \bar{\Phi}'' + 4\bar{\mathcal{D}}\mathcal{D}(J\bar{\Phi}) \end{aligned} \quad (2.1)$$

and

$$\frac{\partial J}{\partial t_3} = J''' + 3([\mathcal{D}, \bar{\mathcal{D}}] J J)' + \frac{3}{2}([\mathcal{D}, \bar{\mathcal{D}}] J^2)' + 4(J^3)' + \frac{3}{2}(\mathcal{D}\bar{\mathcal{D}}\Phi)' + 6(J\Phi\bar{\Phi})',$$

$$\frac{\partial \Phi}{\partial t_3} = \Phi''' + 6\mathcal{D}\bar{\mathcal{D}}\left(\mathcal{D}(J\bar{\mathcal{D}}\Phi) - 2J^2\Phi\right), \quad \frac{\partial \bar{\Phi}}{\partial t_3} = \bar{\Phi}''' - 6\bar{\mathcal{D}}\mathcal{D}\left(\bar{\mathcal{D}}(J\mathcal{D}\bar{\Phi}) - 2J^2\bar{\Phi}\right), \quad (2.2)$$

where prime means differentiation over space coordinate $\{z\}$, and $\mathcal{D}$ and $\bar{\mathcal{D}}$ are the $N = 2$ supersymmetric Grassmann covariant derivatives

$$\mathcal{D} = \frac{\partial}{\partial \theta} - \frac{\bar{\theta}}{2} \frac{\partial}{\partial z}, \quad \bar{\mathcal{D}} = \frac{\partial}{\partial \bar{\theta}} - \frac{\theta}{2} \frac{\partial}{\partial z}, \quad \mathcal{D}^2 = \bar{\mathcal{D}}^2 = 0, \quad \{\mathcal{D}, \bar{\mathcal{D}}\} = -\partial_z \equiv -\partial. \quad (2.3)$$

The superfields Φ and $\bar{\Phi}$ are constrained to be chiral-anti-chiral correspondingly:

$$\mathcal{D}\Phi = \bar{\mathcal{D}}\bar{\Phi} = 0. \quad (2.4)$$

The integrability of the $N = 4$ KdV hierarchy (2.1),(2.2) has been proved in [3, 1] by explicit construction of the Lax operator and the Lax equation

$$L = \mathcal{D}\bar{\mathcal{D}} + \mathcal{D}\bar{\mathcal{D}}\partial^{-1}\left(J + \bar{\Phi}\partial^{-1}\Phi\right)\partial^{-1}\mathcal{D}\bar{\mathcal{D}}, \quad \frac{\partial L}{\partial t_k} = -\left[L_{\geq 1}^k, L\right]. \quad (2.5)$$

where the subscript $\{\geq 1\}$ indicates the strictly differential part of the pseudodifferential operator. One can easily check that the second and third flow equations for the Lax operator (2.5) coincide with (2.1)-(2.2). The infinite set of conserved currents $\mathcal{H}_k$ for the $N = 4$ KdV hierarchy is given by the following expression

$$H_k \equiv \int dz d\theta d\bar{\theta} \mathcal{H}_k = \int dz d\theta d\bar{\theta} \text{Res} \left(L^k \right) \quad (2.6)$$

where the residue is defined as the coefficient before $[\mathcal{D}, \bar{\mathcal{D}}]\partial^{-1}$.

The main attractive feature of the $N = 4$ KdV hierarchy is that it possesses the classical "small" $N = 4$ SCA (with $su(2)$ affine subalgebra) as the second Hamiltonian structure [5] (actually this was the reason why it was called $N = 4$ KdV hierarchy). Indeed, if we define the PB's for the "small" $N = 4$ SCA as

$$\begin{aligned} \{J(1), J(2)\} &= -\left(J\partial + \partial J + \mathcal{D}J\bar{\mathcal{D}} + \bar{\mathcal{D}}J\mathcal{D} + \frac{1}{2}\partial[\mathcal{D}, \bar{\mathcal{D}}]\right)\delta(1, 2), \\ \{J(1), \Phi(2)\} &= (\Phi\bar{\mathcal{D}}\mathcal{D} + \bar{\mathcal{D}}\Phi\mathcal{D})\delta(1, 2), \\ \{J(1), \bar{\Phi}(2)\} &= (\bar{\Phi}\mathcal{D}\bar{\mathcal{D}} + \mathcal{D}\bar{\Phi}\bar{\mathcal{D}})\delta(1, 2), \\ \{\Phi(1), \bar{\Phi}(2)\} &= -2(\partial\mathcal{D}\bar{\mathcal{D}} + 2\mathcal{D}J\bar{\mathcal{D}})\delta(1, 2), \end{aligned} \quad (2.7)$$

where $\delta(1, 2)$ is the $N = 2$ superspace delta-function

$$\delta(1, 2) = (\theta_1 - \theta_2)(\bar{\theta}_1 - \bar{\theta}_2)\delta(z_1 - z_2), \quad (2.8)$$

and the differential operators in the r.h.s are evaluated at the point Z_1 and the derivatives in the r.h.s are assumed to act freely to the right, then the flow equations for the $N = 4$ KdV hierarchy can be written in hamiltonian form as follows

$$\frac{\partial}{\partial t_k} \begin{pmatrix} J \\ \Phi \\ \bar{\Phi} \end{pmatrix} = \left\{ \begin{pmatrix} J \\ \Phi \\ \bar{\Phi} \end{pmatrix}, H_k \right\}. \quad (2.9)$$

The hamiltonians H_2, H_3 which give rise to the second and third flows of $N = 4$ KdV hierarchy (2.1)-(2.2) have the following explicit form

$$\begin{aligned} H_2 &= - \int dz d\theta d\bar{\theta} \left(J^2 - \frac{1}{2} \bar{\Phi} \Phi \right), \\ H_3 &= \int dz d\theta d\bar{\theta} \left([\mathcal{D}, \bar{\mathcal{D}}] J J + \frac{4}{3} J^3 + 2J\Phi\bar{\Phi} + \frac{1}{2} \Phi' \bar{\Phi} \right). \end{aligned} \quad (2.10)$$

2.2 "Fermionic" extensions of N=2, a=4,-2 KdV hierarchies

A close inspection of the Lax operator for the $N = 4$ KdV hierarchy (2.5) gives a hint as to how to construct integrable hierarchies which have $N = 2$ $a = 4$ KdV in the reduction limit $\Phi = \bar{\Phi} = 0$. Indeed, the superfields Φ and $\bar{\Phi}$ appear in the Lax operator only in pairs. So one may wonder whether the Lax equation (2.5) will be consistent if we change the statistics of the superfields $\Phi, \bar{\Phi}$ so that they become fermionic superfields or if we even add an arbitrary number of additional superfields with arbitrary statistics, while keeping the same form of the Lax operator

$$L = \mathcal{D}\bar{\mathcal{D}} + \mathcal{D}\bar{\mathcal{D}}\partial^{-1} \left(J + \sum_i \bar{\Phi}^i \partial^{-1} \Phi^i + \sum_j \bar{\Psi}^j \partial^{-1} \Psi^j \right) \partial^{-1} \mathcal{D}\bar{\mathcal{D}}, \quad (2.11)$$

where $\bar{\Psi}^j, \Psi^j$ are fermionic chiral/anti-chiral superfields. The answer is positive, but the price we have to pay is non-locality of the second Hamiltonian structure [1, 2], even in the case of only one pair of fermionic superfields ($i = 0, j = 1$). Let us note that for this special case the equations of the second and third flows have the same form as their bosonic counterparts (2.1),(2.2).

From the integrability point of view the Lax operator and Lax equation give us the full information about system. However non-locality of the second Hamiltonian structure is a somewhat unappealing property and we would rather prefer to understand the origin of such non-locality. Therefore we rise the question: is it possible to construct some extended system which reproduce these ones as particular cases when some additional fields are set to zero, and are characterized by a *local* second Hamiltonian structure? If such extended systems exist they are certainly more general and more fundamental. In what follows we consider the simplest case with one pair of fermionic chiral-anti-chiral superfields and we will show that it is indeed possible to construct the extended system of the type just described.

The simplest possibility to construct a *local* algebra which contains at least all superfields that feature in the fermionic version of the equations (2.1),(2.2), i.e. the bosonic J together with the fermionic superfields $\Psi, \bar{\Psi}$, is to add one additional pair of chiral-anti-chiral fermionic superfields $H, \bar{H}$. This extended set of superfields forms a closed *local* although *non-linear* algebra

$$\begin{aligned} \{J(1), J(2)\} &= - \left(J\partial + \partial J + \mathcal{D}J\bar{\mathcal{D}} + \bar{\mathcal{D}}J\mathcal{D} + \frac{1}{2} \partial [\mathcal{D}, \bar{\mathcal{D}}] \right) \delta(1, 2), \\ \{J(1), \Psi(2)\} &= \left(2\bar{\mathcal{D}}\mathcal{D}\Psi + \mathcal{D}\Psi\bar{\mathcal{D}} + \bar{\mathcal{D}}\Psi\mathcal{D} - \Psi\partial \right) \delta(1, 2), \\ \{J(1), \bar{\Psi}(2)\} &= \left(2\mathcal{D}\bar{\mathcal{D}}\bar{\Psi} + \mathcal{D}\bar{\Psi}\bar{\mathcal{D}} + \bar{\mathcal{D}}\bar{\Psi}\mathcal{D} - \bar{\Psi}\partial \right) \delta(1, 2), \end{aligned}$$

$$\begin{aligned}
\{J(1), H(2)\} &= H\overline{\mathcal{D}}\mathcal{D}\delta(1,2), \quad \{J(1), \overline{H}(2)\} = \overline{H}\mathcal{D}\overline{\mathcal{D}}\delta(1,2), \\
\{H(1), \overline{H}(2)\} &= \frac{1}{2}\mathcal{D}\overline{\mathcal{D}}\delta(1,2), \quad \{H(1), \Psi(2)\} = \mathcal{D}\Psi\delta(1,2), \quad \{H(1), \overline{\Psi}(2)\} = -\mathcal{D}\overline{\Psi}\delta(1,2), \\
\{\overline{H}(1), \Psi(2)\} &= -\overline{\mathcal{D}}\Psi\delta(1,2), \quad \{\overline{H}(1), \overline{\Psi}(2)\} = \overline{\mathcal{D}}\overline{\Psi}\delta(1,2), \\
\{\Psi(1), \overline{\Psi}(2)\} &= 2\left(-\partial\mathcal{D}\overline{\mathcal{D}} - 2\mathcal{D}J\overline{\mathcal{D}} + \Psi\overline{\Psi} - 4\mathcal{D}J\overline{H} - 4JH\overline{\mathcal{D}} - 8JH\overline{H} + 2\mathcal{D}\overline{H}\mathcal{D}\overline{\mathcal{D}} \right. \\
&\quad - 2\partial\mathcal{D}\overline{H} - 2\partial H\overline{\mathcal{D}} - 2H\partial\overline{\mathcal{D}} + 2\overline{\mathcal{D}}H\mathcal{D}\overline{\mathcal{D}} + 4\mathcal{D}\overline{H}\mathcal{D}\overline{H} + 4H\overline{\mathcal{D}}H\overline{\mathcal{D}} \\
&\quad + 8H\overline{H}\mathcal{D}\overline{H} + 8H\overline{H}\overline{\mathcal{D}}H + 4\mathcal{D}\overline{\mathcal{D}}H\overline{H} + 4H\overline{\mathcal{D}}\mathcal{D}\overline{H} \\
&\quad \left. - 8H\mathcal{D}\overline{\mathcal{D}}\overline{H} + 4H\overline{H}\mathcal{D}\overline{\mathcal{D}}\right)\delta(1,2), \tag{2.12}
\end{aligned}$$

which we claim is the right second Hamiltonian structure for the fermionic version of the system (2.1),(2.2) in the coset approach.

One can check that the Jacobi identities put additional constraints on the superfields $\Psi, \overline{\Psi}$:

$$\mathcal{D}\Psi = -2H\Psi, \quad \overline{\mathcal{D}}\overline{\Psi} = 2\overline{H}\overline{\Psi}. \tag{2.13}$$

The set of Poisson brackets (2.12) together with the covariant chirality constraints (2.13) form the closed algebra we are looking for¹. This algebra has the structure of non-linear W algebra due to the presence of non-linear terms in the right hand side.

The algebra (2.12) possesses manifest $N = 2$ supersymmetry. The same is true also for the coset which is defined by quotienting it by the $\widehat{U(1)}$ subalgebra generated by $H, \overline{H}$. It is this coset that forms the second Hamiltonian structure for the "fermionic" extension of $N = 2, a = 4$ KdV hierarchy. Stated another way, it is possible to construct an infinite number of Hamiltonians which have vanishing Poisson brackets with respect to $H, \overline{H}$. Thus, at the level of equations of motion one can consistently put $H = \overline{H} = 0$ and remain with the equations for the fields $J, \Psi, \overline{\Psi}$. Moreover, after putting H and $\overline{H}$ equal to zero, the covariant chirality conditions (2.13) will become the ordinary chiral/anti-chiral ones

$$\mathcal{D}\Psi = \overline{\mathcal{D}}\overline{\Psi} = 0. \tag{2.14}$$

The Hamiltonian densities $\mathcal{H}_k$ in the $N = 2$ superspace for the integer $k = 1, 2, \dots$ are bosonic and have conformal dimension k . The explicit form of the first three Hamiltonians is

$$\begin{aligned}
H_1 &= \int dzd\theta d\bar{\theta} J, & H_2 &= \int dzd\theta d\bar{\theta} \left(-J^2 + \frac{1}{2}\overline{\Psi}\Psi\right) \\
H_3 &= \int dzd\theta d\bar{\theta} \left(J[\mathcal{D}, \overline{\mathcal{D}}]J + 2J\Psi\overline{\Psi} + \frac{4}{3}J^3 - (\overline{\mathcal{D}}H + \mathcal{D}\overline{H})\Psi\overline{\Psi} + \frac{1}{2}\Psi'\overline{\Psi}\right). \tag{2.15}
\end{aligned}$$

The flows are defined by

$$\frac{\partial}{\partial t_k}\mathcal{A}(Z) = \left\{\mathcal{A}(Z), \int dX\mathcal{H}_k(X)\right\} \tag{2.16}$$

for a given superfield $\mathcal{A}$. Since all the Hamiltonians (2.15) have vanishing Poisson brackets with $H, \overline{H}$, we can consistently put $H = \overline{H} = 0$ in the r.h.s. of (2.16) *after* evaluating the Poisson brackets.

¹To our best knowledge, this algebra first appeared in hidden form (as subalgebra of some extended algebra in a special basis) in the Ref.[6].

The first flow just gives the following equations of motion for $J, \Psi, \bar{\Psi}$

$$\frac{\partial}{\partial t_1} J = J' , \quad \frac{\partial}{\partial t_1} \Psi = \Psi' , \quad \frac{\partial}{\partial t_1} \bar{\Psi} = \bar{\Psi}' , \quad (2.17)$$

while the second and third Hamiltonians generate the equations (2.1), (2.2).

It is interesting that the same coset describes another integrable hierarchy [4] which is an extension of $N = 2$ KdV hierarchy with $a = -2$ [7] and can be defined by the following Lax operator and Lax equation [4]:

$$L = \mathcal{D}\bar{\mathcal{D}}\partial + 2\mathcal{D}J\bar{\mathcal{D}} + \mathcal{D}\Psi\partial^{-1}\bar{\Psi}\bar{\mathcal{D}} , \quad \frac{\partial}{\partial t_k} L = - \left[\left(L^{\frac{k}{2}} \right)_{\geq 1} , L \right] . \quad (2.18)$$

The first two non-trivial (i.e. the second and third) flow equations of motion are

$$\frac{\partial J}{\partial t_2} = -(\bar{\Psi}\Psi)' , \quad \frac{\partial \Psi}{\partial t_2} = \Psi'' - 2\mathcal{D}(J\bar{\mathcal{D}}\Psi) , \quad \frac{\partial \bar{\Psi}}{\partial t_2} = -\bar{\Psi}'' - 2\bar{\mathcal{D}}(J\mathcal{D}\bar{\Psi}) \quad (2.19)$$

and

$$\begin{aligned} \frac{\partial J}{\partial t_3} &= J''' + 3([\mathcal{D}, \bar{\mathcal{D}}] J J)' - \frac{3}{2}([\mathcal{D}, \bar{\mathcal{D}}] J^2)' - 2(J^3)' - 3(\Psi\bar{\Psi}' - \Psi'\bar{\Psi})' + 6(J\Psi\bar{\Psi})' , \\ \frac{\partial \Psi}{\partial t_3} &= 4\Psi''' - 6\mathcal{D}(\bar{\mathcal{D}}\Psi J' + 2\bar{\mathcal{D}}\Psi' J + J^2\bar{\mathcal{D}}\Psi + \Psi\bar{\Psi}\bar{\mathcal{D}}\Psi) , \\ \frac{\partial \bar{\Psi}}{\partial t_3} &= 4\bar{\Psi}''' + 6\bar{\mathcal{D}}(\mathcal{D}\bar{\Psi} J' + 2\mathcal{D}\bar{\Psi}' J - J^2\mathcal{D}\bar{\Psi} - \Psi\bar{\Psi}\mathcal{D}\bar{\Psi}) . \end{aligned} \quad (2.20)$$

Like the previous case the algebra (2.12) provides the second Hamiltonian structure for the hierarchy (2.18) via the coset construction. Explicitly, the first three Hamiltonians read

$$\begin{aligned} H_1 &= \int dz d\theta d\bar{\theta} J , \quad H_2 = \int dz d\theta d\bar{\theta} \left(\frac{1}{2} \Psi \bar{\Psi} \right) , \\ H_3 &= \int dz d\theta d\bar{\theta} \left(J [\mathcal{D}, \bar{\mathcal{D}}] J + 2\Psi'\bar{\Psi} + 2J\Psi\bar{\Psi} - \frac{2}{3}J^3 - 4(\bar{\mathcal{D}}H + \mathcal{D}\bar{H})\Psi\bar{\Psi} \right) , \end{aligned} \quad (2.21)$$

where Ψ and $\bar{\Psi}$ are subject to the constraints (2.13) and the different flow equations are defined by (2.16).

Thus both, "fermionic" extensions of $N = 2$ KdV hierarchy with $a = 4$ and $a = -2$ can be treated within the coset approach as $N = 2$ quotients of the $N = 2$ nonlinear algebra (2.12). It is remarkable that although the other details of these two hierarchies are different, the second hamiltonian structure is the same, as was pointed out above.

3 Second Hamiltonian structure

In the previous section we used the coset formalism applied to the $N = 2$ algebra (2.12) to describe the "fermionic" extensions of $N = 2$ $a = 4, -2$ KdV hierarchies. This approach allows for a simpler analysis than the Dirac constraint procedure, because the constraints $H = \bar{H} = 0$ can be imposed in the equations of motion. In this Section we will show how

to apply the constraints $H = \overline{H} = 0$ to the algebra (2.12) and will demonstrate that the Dirac brackets for the remaining supercurrents $J, \Psi, \overline{\Psi}$ close on a non-local algebra [1, 2].

Let us start with the obvious remark that the constraints

$$H = \overline{H} = 0, \quad (3.1)$$

we want to apply, are incompatible with the Poisson brackets of our algebra

$$\{H(1), \overline{H}(2)\} = \frac{1}{2} \mathcal{D} \overline{\mathcal{D}} \delta(1, 2), \quad \{\overline{H}(1), H(2)\} = -\frac{1}{2} \overline{\mathcal{D}} \mathcal{D} \delta(1, 2), \quad (3.2)$$

so we need to apply Dirac's procedure passing to the Dirac brackets

$$\{\mathcal{A}, \mathcal{B}\}^* \equiv \{\mathcal{A}, \mathcal{B}\} - \{\mathcal{A}, H\} \left(-\frac{1}{2} \overline{\mathcal{D}} \mathcal{D}\right)^{-1} \{\overline{H}, \mathcal{B}\} - \{\mathcal{A}, \overline{H}\} \left(\frac{1}{2} \mathcal{D} \overline{\mathcal{D}}\right)^{-1} \{H, \mathcal{B}\} \quad (3.3)$$

for any two superfields $\mathcal{A}, \mathcal{B}$. One can expect some extra complications in the Dirac procedure because the operators $\mathcal{D} \overline{\mathcal{D}}$ and $\overline{\mathcal{D}} \mathcal{D}$ are not invertible on the whole $N = 2$ superspace. But fortunately this is not so for the case at hand. Indeed, due to the chiral-anti-chiral structure of H and $\overline{H}$, the Poisson brackets $\{\mathcal{A}, H\}, \{\overline{H}, \mathcal{B}\}, \{\mathcal{A}, \overline{H}\}, \{H, \mathcal{B}\}$ can be always rewritten in the following form

$$\begin{aligned} \{\mathcal{A}, H\} &\sim (\dots) \mathcal{D} \delta(1, 2), & \{\mathcal{A}, \overline{H}\} &\sim (\dots) \overline{\mathcal{D}} \delta(1, 2), \\ \{H, \mathcal{B}\} &\sim \mathcal{D} (\dots) \delta(1, 2), & \{\overline{H}, \mathcal{B}\} &\sim \overline{\mathcal{D}} (\dots) \delta(1, 2), \end{aligned} \quad (3.4)$$

where $(\dots)$ is used to denote any combinations of operators and/or superfields. Thus, one can immediately conclude that we have to know the operators $(\mathcal{D} \overline{\mathcal{D}})^{-1}$ and $(\overline{\mathcal{D}} \mathcal{D})^{-1}$ not on the whole $N = 2$ superspace but only on its chiral-anti-chiral subspaces, where they are well defined. Explicitly

$$\begin{aligned} \mathcal{D} \left(-\frac{1}{2} \overline{\mathcal{D}} \mathcal{D}\right)^{-1} \overline{\mathcal{D}} &= \mathcal{D} (2\partial^{-1}) \overline{\mathcal{D}} \quad \text{because} \quad \mathcal{D} (-\overline{\mathcal{D}} \mathcal{D} \partial^{-1}) \overline{\mathcal{D}} \equiv \mathcal{D} \overline{\mathcal{D}}, \\ \overline{\mathcal{D}} \left(\frac{1}{2} \mathcal{D} \overline{\mathcal{D}}\right)^{-1} \mathcal{D} &= \overline{\mathcal{D}} (-2\partial^{-1}) \mathcal{D} \quad \text{because} \quad \overline{\mathcal{D}} (-\mathcal{D} \overline{\mathcal{D}} \partial^{-1}) \mathcal{D} \equiv \overline{\mathcal{D}} \mathcal{D}. \end{aligned} \quad (3.5)$$

Keeping this in the mind all calculations can be done straightforwardly.

All Dirac brackets containing $H, \overline{H}$ are now equal to zero, so we can impose the constraints (3.1) in strong sense. The rest of the Dirac brackets between the superfields $J, \Psi, \overline{\Psi}$ form the following closed non-local algebra

$$\begin{aligned} \{J(1), J(2)\}^* &= -\left(J\partial + \partial J + \mathcal{D} J \overline{\mathcal{D}} + \overline{\mathcal{D}} J \mathcal{D} + \frac{1}{2} \partial [\mathcal{D}, \overline{\mathcal{D}}]\right) \delta(1, 2), \\ \{J(1), \Psi(2)\}^* &= (-\overline{\mathcal{D}} \Psi \mathcal{D} + \Psi [\mathcal{D}, \overline{\mathcal{D}}]) \delta(1, 2), \\ \{J(1), \overline{\Psi}(2)\}^* &= (-\mathcal{D} \overline{\Psi} \overline{\mathcal{D}} + \overline{\Psi} [\mathcal{D}, \overline{\mathcal{D}}]) \delta(1, 2), \\ \{\Psi(1), \Psi(2)\}^* &= -4\Psi \overline{\mathcal{D}} \mathcal{D} \partial^{-1} \Psi \delta(1, 2), \quad \{\overline{\Psi}(1), \overline{\Psi}(2)\}^* = 4\overline{\Psi} \mathcal{D} \overline{\mathcal{D}} \partial^{-1} \overline{\Psi} \delta(1, 2), \\ \{\Psi(1), \overline{\Psi}(2)\}^* &= -2(\mathcal{D} \overline{\mathcal{D}} \partial + 2\mathcal{D} J \overline{\mathcal{D}} - 2\Psi \mathcal{D} \overline{\mathcal{D}} \partial^{-1} \overline{\Psi}) \delta(1, 2) \end{aligned} \quad (3.6)$$

with $\Psi, \bar{\Psi}$ obeying the chirality constraints

$$\mathcal{D}\Psi = \bar{\mathcal{D}}\bar{\Psi} = 0. \quad (3.7)$$

The algebra (3.6) is a particular case of the non-local algebras proposed in [1, 2]. Let us note that validity of the Jacobi identities for the non-local algebra (3.6) is guaranteed since we constructed it by applying the standard Dirac's procedure applied to the local algebra (2.12).

Now one can straightforwardly check that the Hamiltonians

$$\begin{aligned} H_1 &= \int dz d\theta d\bar{\theta} J, & H_2 &= \int dz d\theta d\bar{\theta} \left(-J^2 + \frac{1}{2} \bar{\Psi} \Psi \right), \\ H_3 &= \int dz d\theta d\bar{\theta} \left(J [\mathcal{D}, \bar{\mathcal{D}}] J + 2J\Psi\bar{\Psi} + \frac{4}{3} J^3 + \frac{1}{2} \Psi' \bar{\Psi} \right) \end{aligned} \quad (3.8)$$

and

$$\begin{aligned} H_1 &= \int dz d\theta d\bar{\theta} J, & H_2 &= \int dz d\theta d\bar{\theta} \left(\frac{1}{2} \Psi \bar{\Psi} \right), \\ H_3 &= \int dz d\theta d\bar{\theta} \left(J [\mathcal{D}, \bar{\mathcal{D}}] J + 2\Psi' \bar{\Psi} + 2J\Psi\bar{\Psi} - \frac{2}{3} J^3 \right) \end{aligned} \quad (3.9)$$

produce the correct flow equations for the "fermionic" extensions of $N = 2$ $a = 4$ and $a = -2$ KdV hierarchies; they thus correspondingly provide the second Hamiltonian structure for both hierarchies.

4 New extensions of N=2 KdV

In the previous sections we have showed that the second hamiltonian structure for the "fermionic" extensions of the $N = 2$ $a = 4$ and $a = -2$ KdV hierarchies in the coset approach is given by the very non trivial non-local algebra (3.6). At the same time these hierarchies can be obtained via hamiltonian reduction starting from the local algebra (2.12). One may wonder whether there exist non trivial integrable hierarchies that include all the superfields $(J, \Psi, \bar{\Psi}, H, \bar{H})$ and possess the algebra (2.12) as their second Hamiltonian structure. This question is rather complicated and will be addressed elsewhere. Here, instead, we will discuss a simpler problem concerning the existence of integrable hierarchies with second Hamiltonian structure given by the following *linear* subalgebra of (2.12)

$$\begin{aligned} \{J(1), J(2)\} &= - \left(J\partial + \partial J + \mathcal{D}J\bar{\mathcal{D}} + \bar{\mathcal{D}}J\mathcal{D} - \frac{1}{2} \partial [\mathcal{D}, \bar{\mathcal{D}}] \right) \delta(1, 2), \\ \{H(1), \bar{H}(2)\} &= \mathcal{D}\bar{\mathcal{D}}\delta(1, 2), \end{aligned} \quad (4.1)$$

while all the other PB's vanish. This is the direct product of the $N = 2$ SCA and the affine $\hat{U}(1)$ algebra. We will show that there exist at least the first three non-trivial flows, or, in other words the first four Hamiltonians ($H_1 - H_4$) which have the proper dimensions and are mutually commuting with respect to the Poisson brackets (4.1). Despite the very trivial Hamiltonian structure there are three non-equivalent systems for which we

succeeded in finding the first four hamiltonians. The first Hamiltonian is the same for all cases $H_1 = \int dz d\theta \bar{\theta} (J + H\bar{H})$ and corresponds to the trivial flow equations, like (2.17). Due to the very complicated structure of H_3 and H_4 , we present here, in addition, only the second hamiltonians for these systems together with the corresponding second flow equations (for the explicit form of the third Hamiltonians see Appendix).

Extension of $N = 2, a = 4$ KdV. The first system is given by the following Hamiltonian density and second flow equations

$$\mathcal{H}_2^{(1)} = J^2 + H\bar{H}' + 4JH\bar{H} + \sqrt{2}H\bar{H}(\bar{\mathcal{D}}H + \mathcal{D}\bar{H}) \quad (4.2)$$

$$\begin{aligned} \frac{\partial J}{\partial t_2} &= -[\mathcal{D}, \bar{\mathcal{D}}] J' - 4JJ' + 4\bar{\mathcal{D}}J\mathcal{D}(H\bar{H}) + 4\mathcal{D}J\bar{\mathcal{D}}(H\bar{H}) \\ &\quad + (4JH\bar{H} - 4\bar{\mathcal{D}}H\mathcal{D}\bar{H} + H'\bar{H} - H\bar{H}')', \\ \frac{\partial H}{\partial t_2} &= -H'' - \mathcal{D} \left(4\bar{\mathcal{D}}(JH) + \sqrt{2}\bar{\mathcal{D}}H\mathcal{D}\bar{H} + 2\sqrt{2}(H\bar{H})' \right), \\ \frac{\partial \bar{H}}{\partial t_2} &= \bar{H}'' - \bar{\mathcal{D}} \left(4\mathcal{D}(J\bar{H}) + \sqrt{2}\mathcal{D}\bar{H}\mathcal{D}\bar{H} - 2\sqrt{2}(H\bar{H})' \right). \end{aligned} \quad (4.3)$$

One can immediately see that the system (4.3) admits the reduction $H = \bar{H} = 0$. The latter coincides with the $N = 2, a = 4$ KdV hierarchy.

Extension of $N = 2, a = -2$ KdV. The second solution one can construct is described by the following flow equations

$$\begin{aligned} \frac{\partial J}{\partial t_2} &= \bar{\mathcal{D}}J\mathcal{D}(H\bar{H}) + \mathcal{D}J\bar{\mathcal{D}}(H\bar{H}) + \left(JH\bar{H} - \bar{\mathcal{D}}H\mathcal{D}\bar{H} + \frac{1}{2}(H'\bar{H} - H\bar{H}') \right)', \\ \frac{\partial H}{\partial t_2} &= \frac{1}{2}H'' - \mathcal{D} \left(\bar{\mathcal{D}}(JH) + \frac{\sqrt{2}}{2}\bar{\mathcal{D}}H\mathcal{D}\bar{H} + \sqrt{2}(H\bar{H})' \right), \\ \frac{\partial \bar{H}}{\partial t_2} &= -\frac{1}{2}\bar{H}'' - \bar{\mathcal{D}} \left(\mathcal{D}(J\bar{H}) + \frac{\sqrt{2}}{2}\mathcal{D}\bar{H}\mathcal{D}\bar{H} - \sqrt{2}(H\bar{H})' \right), \end{aligned} \quad (4.4)$$

with the Hamiltonian density

$$\mathcal{H}_2^{(2)} = -\frac{1}{2}H\bar{H}' + JH\bar{H} + \frac{1}{\sqrt{2}}H\bar{H}(\bar{\mathcal{D}}H + \mathcal{D}\bar{H}) \quad (4.5)$$

Again the equations (4.4) admit the solution $H = \bar{H} = 0$, which corresponds to the well-known $N = 2, a = -2$ KdV hierarchy.

New $N = 2$ supersymmetric system. It comes as a surprise that there is a third solution, which does not correspond to any known integrable system. It is described by the following Hamiltonian density and corresponding second flow equations:

$$\mathcal{H}_2^{(3)} = J^2 - 3H\bar{H}' + 2\sqrt{2}J(\bar{\mathcal{D}}H + \mathcal{D}\bar{H}) + \sqrt{2}H\bar{H}(\bar{\mathcal{D}}H + \mathcal{D}\bar{H}) \quad (4.6)$$

and

$$\begin{aligned}
\frac{\partial J}{\partial t_2} &= -[\mathcal{D}, \overline{\mathcal{D}}] J' + 4JJ' - 2\sqrt{2}(\overline{\mathcal{D}}JH' + \mathcal{D}J\overline{H}') \\
&\quad + \sqrt{2}(\mathcal{D}\overline{H}' - \overline{\mathcal{D}}H' + 2J\overline{\mathcal{D}}H + 2J\mathcal{D}\overline{H})', \\
\frac{\partial H}{\partial t_2} &= 3H'' - 2\sqrt{2}\mathcal{D}\left(J' + \frac{1}{2}\overline{\mathcal{D}}H\overline{\mathcal{D}}H + (H\overline{H})'\right), \\
\frac{\partial \overline{H}}{\partial t_2} &= -3\overline{H}'' + 2\sqrt{2}\overline{\mathcal{D}}\left(J' - \frac{1}{2}\mathcal{D}\overline{H}\mathcal{D}\overline{H} + (H\overline{H})'\right). \tag{4.7}
\end{aligned}$$

One can see that the system (4.7) does not admit the solution $H = \overline{H} = 0$ and so it cannot be reduced, like the previous cases, to a KdV-type equation on a single superfield J .

Of course, the existence of the first four Hamiltonians for the three systems does not guarantee by itself their integrability but it is a good reason to take these systems seriously (i.e. search for Lax operators and Lax equations).

Finally let us stress, that if we take, instead of the Hamiltonian structure (4.1), the direct product of two $N = 2$ SCA's (e.g. constructing the second $N = 2$ SCA through the Sugawara procedure from $H, \overline{H}$) we do not succeed in finding any system admitting second flows [8].

5 Discussions and outlook

In this letter we have described the two hierarchies - the "fermionic" extensions of $N = 2$ KdV with $a = 4$ (2.15) and $a = -2$ (2.21) within the coset framework. We have explicitly showed that both hierarchies can be obtained by quotienting the $N = 2$ superalgebra (2.12) by its $N = 2$ $\widehat{U(1)}$ subalgebra. We also applied Dirac's procedure to construct the second Hamiltonian structure for these hierarchies. The resulting $N = 2$ superalgebra (3.6) is non-local for these hierarchies in contrast with the former $N = 4$ KdV [5] and "bosonic" extension of the $N = 2$ KdV hierarchy [4], which possess local linear "small" $N = 4$ SCA as second Hamiltonian structure. We also proposed three additional (presumably) integrable extensions of the $N = 2$ KdV hierarchy which possess at least the four first conserved currents.

The main advantage of the coset approach is that the original $N = 2$ superalgebra (2.12) is *local*. One may therefore conjecture that also other hierarchies which can be obtained from the Lax operator (2.11) can also be described as a coset of some *local* $N = 2$ superalgebra. Moreover, as was shown in [9] for the NLS case, these extended superalgebras, which localized the second Hamiltonian structure for NLS hierarchy [10], could give rise to new hierarchies with an extended set of superfields. We defer to a future publication the analysis of whether it is possible to construct new integrable hierarchies with the superalgebra (2.12) as second Hamiltonian structure.

Another important comment concerns the possible existence of other $N = 2$ GNLS hierarchies, besides those constructed in [11]. Indeed, while the known $N = 2$ GNLS hierarchy corresponds to the $N = 2$ $a = 4$ KdV and its bosonic/fermionic extensions, the existence of the "bosonic" and/or "fermionic" extensions of $N = 2$ KdV hierarchies [4]

with $a = -2$ with the same second Hamiltonian structure seems to imply existence of the corresponding "new" GNLS hierarchies. Such "new" $N = 2$ GNLS hierarchies do exist, and will be considered in a forthcoming paper [12].

Acknowledgments. S.K. is grateful to E. Ivanov for many useful discussions. He also thanks SISSA for the hospitality and financial support extended to him during the course of this work.

This investigation has been supported in part by the Russian Foundation of Fundamental Research, grant No. 96-02-17634, RFBR-DFG grant No. 96-02-00180, INTAS grant 94-2317, and by a grant from the Dutch NWO organization.

Appendix

Here we write down the expressions for the third Hamiltonians for the systems (4.3), (4.4), (4.7). The general expression for the third Hamiltonian density have the following form

$$\begin{aligned} \mathcal{H}_3 = & a_1([\mathcal{D}, \overline{\mathcal{D}}] J)J + a_2 J \overline{\mathcal{D}} H' + a_3 J \mathcal{D} \overline{H}' + a_4 H \overline{H}'' + a_5 (\overline{\mathcal{D}} H') H \overline{H} + a_6 ([\mathcal{D}, \overline{\mathcal{D}}] J) H \overline{H} \\ & + a_7 H (\mathcal{D} \overline{H}') \overline{H} + a_8 (\overline{\mathcal{D}} J) (\overline{\mathcal{D}} H) H + a_9 (\overline{\mathcal{D}} J) J H + a_{10} (\mathcal{D} J) (\mathcal{D} \overline{H}) \overline{H} + a_{11} (\mathcal{D} J) J \overline{H} \\ & + a_{12} J (\overline{\mathcal{D}} H) (\mathcal{D} \overline{H}) + a_{13} J^3 + a_{14} (\overline{\mathcal{D}} H)^2 H \overline{H} + a_{15} (\overline{\mathcal{D}} H) H (\mathcal{D} \overline{H}) \overline{H} \\ & + a_{16} (\mathcal{D} \overline{H})^2 H \overline{H} + a_{17} J (\overline{\mathcal{D}} H) H \overline{H} + a_{18} J (\mathcal{D} \overline{H}) H \overline{H} + a_{19} J^2 H \overline{H}. \end{aligned} \quad (\text{A.1})$$

The solutions we obtained are

Case	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9
a=4 KdV	1	0	0	-1	$3\sqrt{2}$	0	$-3\sqrt{2}$	0	0
a=-2 KdV	-1/6	0	0	2/3	$2\sqrt{2}$	-1	$-2\sqrt{2}$	0	0
new KdV	1/3	$1/\sqrt{2}$	$-1/\sqrt{2}$	2/3	$3/(8\sqrt{2})$	1	$-3/\sqrt{2}$	-1	$-\sqrt{2}$
a_{10}	a_{11}	a_{12}	a_{13}	a_{14}	a_{15}	a_{16}	a_{17}	a_{18}	a_{19}
0	0	6	-4/3	-2	0	-2	$-6\sqrt{2}$	$-6\sqrt{2}$	-12
0	0	2	-1/9	4/3	3	4/3	$2\sqrt{2}$	$2\sqrt{2}$	1
-1	$-\sqrt{2}$	1	2/9	1/3	1	1/3	0	0	0

References

- [1] F. Delduc and L. Gallot, " $N=2$ KP and KdV hierarchies in extended superspace", ENSLAPP-L-617/96, solv-int/9609008.
- [2] L. Bonora and A. Sorin, " *The Hamiltonian structure of the $N=2$ supersymmetric GNLS hierarchy*", SISSA 52/97/EP, hep-th/9704130.
- [3] E. Ivanov and S. Krivonos, " *New integrable extensions of $N=2$ KdV and Boussinesq hierarchies*", JINR E2-96-344, hep-th/9609191, (Phys. Lett. A in press).

- [4] F. Delduc, L. Gallot and E. Ivanov, Phys. Lett. **B 396** (1997) 122.
- [5] F. Delduc and E. Ivanov, Phys. Lett. **B 309** (1993) 312.
- [6] C. Ahn, E. Ivanov and A. Sorin, Commun. Math. Phys. **183** (1997) 205.
- [7] P. Labelle and P. Mathieu, J. Math. Phys. **32** (1991) 923.
- [8] Z. Popowicz, “*Integrable system constructed out of two interacting superconformal fields*”, hep-th/9702090.
- [9] E. Ivanov, S. Krivonos and F. Toppan, “ *$N=4$ Super NLS-mKdV Hierarchies*”, DFPD 97-TH-11, E2-97-108, hep-th/9703224.
- [10] S. Krivonos, A. Sorin and F. Toppan, Phys. Lett. **A 206** (1995) 146.
- [11] L. Bonora, S. Krivonos and A. Sorin, Nucl. Phys. **B 477** (1996) 835.
- [12] L. Bonora and S. Krivonos, “*New $N=2$ GNLS hierarchy*”, in preparation.